


# TROPOSPHERIC OZONE FROM IASI: COMPARISON OF DIFFERENT INVERSION ALGORITHMS AND VALIDATION WITH OZONE SONDES

A. Griesfeller<sup>1</sup>, C. Keim<sup>2</sup>, G. Dufour<sup>1</sup>, M. Eremenko<sup>1</sup>, J.-M. Flaud<sup>1</sup>, J. Orphal<sup>3</sup>, M. Höpfner<sup>3</sup>, C. Clerbaux<sup>4</sup>, C. Scannell<sup>4</sup>, P.-F. Coheur<sup>5</sup>, D. Hurtmans<sup>5</sup>, S. Payan<sup>6</sup>, B. Barret<sup>7</sup>, E. Le Flochmoën<sup>7</sup>

1 LISA, CNRS, France, 2 Astrium GmbH, Germany , 3 IMK-ASF, KIT, Germany, 4 LATMOS-IPSL, France, 5 ULB, Belgium, 6 LPMAA, France, 7 LA CNRS, France


## First comparison results:

Comparison of the inversion algorithms from LISA, LATMOS, LPMAA, and Eumesat:  
June 2007-August 2008



**Comparison of the inversion algorithms from LISA and LA for 3 months: June 2008 – August 2008:  
50 coincidences for 8 stations in the NH midlatitudes**

**Tropospheric profiles**



**Partial column amounts**

